

## TOLFINE

### Injectable solution

#### COMPOSITION

Tolfenamic acid	0,0400
Benzyl alcohol (E1519)	0,0104
Sodium hydroxymethanesulfinate	0,0050
Excipient QS	1 ml

#### INDICATIONS

In cattle:

- Reduction of inflammation associated with musculoskeletal and respiratory infections

#### CONTRAINDICATIONS

The use of this veterinary medicine is not recommended in the case of heart, liver or kidney failure.

Tolfenamic acid is contraindicated in the case of ulcerations or digestive bleeding.

Do not use in case of hypersensitivity to the active substances or to any of the excipients.

#### ADVERSE REACTIONS

A transient inflammation and tumefaction may appear at the injection site. If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon

#### TARGET SPECIES

Cattle

## **DOSAGE, ROUTE AND METHOD OF ADMINISTRATION**

2 mg per kg bodyweight, i.e. 1 ml per 20 kg bodyweight, by intramuscular injection, to be repeated every 48 hours.

2 mg per kg bodyweight, i.e. 1 ml per 20 kg bodyweight, by intravenous injection, to be repeated every 24 hours, or to be relayed 24 hours later with intramuscular injections.

Do not exceed 5 consecutive injections

## **WITHDRAWAL PERIOD**

Intramuscular route:

Meat and offal: 12 days. Milk: zero day

Intravenous route:

Meat and offal: 4 days. Milk: 12 hours

## **SPECIAL STORAGE PRECAUTIONS**

After opening: 28 days.

Protect from light.

Keep out of the sight and reach of children.

Do not store above 25°C.

## **SPECIAL WARNINGS**

### **Special warnings for each target species**

Do not inject more than 20 ml per injection site.

### **Special precautions for use in animals**

Use aseptic precautions when administering the product.

Do not use in dehydrated, hypovolaemic or hypotensive animals

(potential risk of increased renal toxicity).

The concurrent administration of potentially nephrotoxic drugs must be avoided

**Special precautions to be taken by the person administering the medicinal product to animals**

In case of contact with the skin or eyes, immediately rinse with water

**Use during pregnancy or lactation**

The safety of the product has not been established during pregnancy

Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

**Interaction with other medicinal products and other forms of interaction**

Do not administer other non-steroidal anti-inflammatories concurrently or within 24 hours of each other. Tolfenamic acid is strongly bound to plasma proteins and may compete with other strongly bound drugs

**Overdose**

At high doses, transient neurological disorders have been observed in cows.

**SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local regulations.

**PRESENTATIONS**

10 ml vial, 25 ml vial, 30 ml vial, 50 ml vial, 100 ml vial, 250 ml vial

**FOR ANIMAL TREATMENT ONLY**  
**TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION**