

Selepherol

COMPOSITION

Selenium(as sodium selenite)	0.023g
VitaminE(as acetate)	3.820g
Sorbic acid	0.050g
Methylparahydroxybenzoate(E218)	0.10g
Propylparahydroxybenzoate(E216)	0.02g
Excipient(containing vitamin B12)	to100ml

PRESENTATION:

Vial of 100ml

PROPERTIES

Pharmacological properties

Pharmacotherapeutic group:Alimentary tract and metabolism,seleniumcombinations.

INDICATIONS

In foals,calves,lambs,prevention and treatment of selenium and vitamin E deficiencies:

–Myopathies

–Musculardystrophies.

CONTRA-INDICATIONS

Do not use in cases of known hypersensitivity to active substance or to any of the excipients.

Do not administer to adult horses.

DOSAGE AND ADMINISTRATION

For deep and strict I.M. route. Prevention (only one administration) : Calves, foals:1.5mlper10kg of body weight until100kg, 1mlper10Kg above 100kg of body weight. (refer below : special warnings) . Lambs:0.2ml per kg of body weight.

Therapy: The same dosage as above, repeating injections at 2 or 3 week-intervals if necessary. Shake well before use.

SPECIALWARNINGS

Special precautions for use in animals

Due to the toxicity of selenium , the doses and delays mentioned in the section “Dosage and administration” should be respected .The injections should be performed under rigorous asepsis.

Special precautions to be taken by the person administering the veterinary medicinal product to animals.

In the case of accidental self-injection,seek medical advice immediately

And show the package leaflet to the physician.

Use during pregnancy and lactation

The safety of this medicinal product has not been studied during pregnancy . However, its use during pregnancy does not seem to pose any particular problem.

Interaction with other medicaments and other forms of interaction

No known interaction.

Overdose

Overdose of selenium may induce fast and thread pulse ,dyspnea, colic, polyuria, cyanosis , prostration, depression and ataxia. Incase of overdose, administer symptomatic treatment associated to a correct hydration (infusionor ad libitum watering).

ADVERSEREACTIONS

Inflammatory reaction at the injection site may occur in very rare cases. Reactions such as anaphylactic shock with cardiovascular disorders, prostration, coma possibly leading to the death of the animal have also been observed in very rare cases .

WITHDRAWALPERIODS

Meat and offal:4days

SPECIAL STORAGE PRECAUTIONS

Shelf-life after first opening of the container:28days

Keep out of the reach and sight of children.

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS

Dispose of any unused product and empty containers inaccordance with guidance from your local waste regulation authority.

FOR ANIMAL TREATMENT ONLY

TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION