

Septotryl Injectable

Anti-Infectious

COMPOSITION

Trimethoprim	4g
Sulfamethoxypyridazine	20g
Benzylalcohol	1ml
Excipient	100ml

PRESENTATIONS

Vial of 100ml-Vial of 250ml

TARGET SPECIES: CATTLE-SHEEP

Slow I.V, I.M, S.C. administration.

INDICATIONS

Infections sensitive to the combination of Trimethoprim and Sulfamethoxypyridazine in cattle and sheep.

Treatment of respiratory, gastrointestinal and urinary infections caused by bacteria sensitive to Sulphamethoxypyridazine and trimethoprim.

DOSAGE

CATTLE-SHEEP

Slow I.V. ,I.M. ,S.C. administration

13.4 mg sulphamethoxypyridazine and 2.7 mg trimethoprim per kg b.w. per day by slow I.V. or I.M. route,i.e.1 ml/15kg b.w.

In cattle:

-60 kg calves:4ml

- adults : 30ml

In sheep:

-Lambs: 0.1ml/1.5kg b.w.

-adults: 4ml

CONTRA-INDICATIONS

Do not use in the following cases: Animals with severe renal or hepatic failure.
Dehydration.
Pregnancy.
Newborns.
Hypersensitivity to one of the active ingredients or one of the excipients.
Do not use in horses.

SIDE-EFFECTS

If treatment is prolonged , the synthesis of thyroid hormones may be inhibited.

SPECIAL PRECAUTIONS FOR USE

Do not mix SEPTOTRYL® with other drugs in the same syringe.
Inject slowly without mixing blood in the syringe containing SEPTOTRYL®.

STORAGE CONDITIONS

Shelf-life after first opening : 28days .Do not store above25°C.
Keep out of the reach and sight of children.

WITHDRAWAL TIMES

Meat and offals : 12days.
Milk: 6days (12milking).

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

FOR ANIMAL TREATMENT ONLY

TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION